

Journal Pre-proof

A MULTINATIONAL REPORT TO CHARACTERISE SARS-CoV-2 INFECTION IN PEOPLE WITH CYSTIC FIBROSIS

Rebecca Cosgriff , Susannah Ahern , Scott C. Bell , Keith Brownlee , Pierre-Régis Burgel , Cass Byrnes , Harriet Corvol , Stephanie Y. Cheng , Alexander Elbert , Albert Faro , Christopher H. Goss , Vincent Gulmans , Bruce C. Marshall , Edward McKone , Peter G. Middleton , Rasa Ruseckaite , Anne L. Stephenson , Siobhán B Carr



PII: S1569-1993(20)30125-9
DOI: <https://doi.org/10.1016/j.jcf.2020.04.012>
Reference: JCF 2029

To appear in: *Journal of Cystic Fibrosis*

Received date: 20 April 2020
Revised date: 22 April 2020
Accepted date: 22 April 2020

Please cite this article as: Rebecca Cosgriff , Susannah Ahern , Scott C. Bell , Keith Brownlee , Pierre-Régis Burgel , Cass Byrnes , Harriet Corvol , Stephanie Y. Cheng , Alexander Elbert , Albert Faro , Christopher H. Goss , Vincent Gulmans , Bruce C. Marshall , Edward McKone , Peter G. Middleton , Rasa Ruseckaite , Anne L. Stephenson , Siobhán B Carr , A MULTINATIONAL REPORT TO CHARACTERISE SARS-CoV-2 INFECTION IN PEOPLE WITH CYSTIC FIBROSIS, *Journal of Cystic Fibrosis* (2020), doi: <https://doi.org/10.1016/j.jcf.2020.04.012>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V. on behalf of European Cystic Fibrosis Society.

Highlights

- A report on the outcomes of 40 people with CF who were positive for SARS-CoV-2
- The reported cohort is heterogeneous and includes 11 lung transplant patients
- The data was collected through the Registries of the 8 participating countries
- The clinical course of SARS-CoV-2 in CF appears similar to the general population
- The outcomes of this early cases have been better than initially predicted

Journal Pre-proof

A MULTINATIONAL REPORT TO CHARACTERISE SARS-CoV-2 INFECTION IN PEOPLE WITH CYSTIC FIBROSIS

Authors

Rebecca Cosgriff^a, Susannah Ahern^b, Scott C. Bell^c, Keith Brownlee^a, Pierre-Régis Burgel^d, Cass Byrnes^e, Harriet Corvol^f, Stephanie Y. Cheng^g, Alexander Elbert^h, Albert Faro^h, Christopher H. Gossⁱ, Vincent Gulmans^j, Bruce C. Marshall^h, Edward McKone^k, Peter G. Middleton^l, Rasa Ruseckaite^b, Anne L. Stephenson^m, Siobhán B Carrⁿ

^a Cystic Fibrosis Trust, London, UK; ^b Monash University, Melbourne, Australia; ^c The Prince Charles Hospital, Brisbane, Australia; ^d Université de Paris, Inserm U1016, Institut Cochin and Cochin Hospital, Assistance Publique Hôpitaux de Paris (APHP); ^e University of Auckland, Auckland, New Zealand; ^f Trousseau Hospital, Sorbonne Université, Paris, France; ^g Cystic Fibrosis Canada, Toronto, Canada; ^h Cystic Fibrosis Foundation, Bethesda, US; ⁱ University of Washington, Seattle, US; ^j Dutch CF Foundation NCFs, Baarn, Netherlands; ^k St Vincent's University Hospital, Dublin, Ireland; ^l Westmead Hospital, Sydney, Australia; ^m Toronto Adult Cystic Fibrosis Centre, St. Michael's Hospital, Toronto, Canada; ⁿ Royal Brompton Hospital and Imperial College London, UK

Correspondence

Rebecca Cosgriff, Cystic Fibrosis Trust, 1 Aldgate, London, EC3N 1RE, UK.

Email: rebecca.cosgriff@cysticfibrosis.org.uk

Collaborating authors

Name	Institution	Country
David Reid	The Prince Charles Hospital, Brisbane	Australia
Peter Wark	John Hunter Hospital, Newcastle	Australia
Géraldine Daneau	Institute of Tropical Medicine	Belgium
Véronique Boussaud	Cochin Hospital, Paris	France
Graziella Brinchault	Rennes	France
Emmanuelle Coirier-Duet	Versailles	France
Jean-Christophe Dubus	Marseille	France
Dominique Grenet	Foch Hospital, Suresnes	France
Sandra de Miranda	Foch Hospital, Suresnes	France
Laurence Beaumont	Foch Hospital, Suresnes	France
Reem Kanaan	Cochin Hospital, Paris	France
Muriel Lauraens	Caen	France
Clémence Martin	Cochin Hospital, Paris	France
Marie Mittaine	Toulouse	France
Anne Prévotat	Lille	France
Martine Reynaud-Gaubert	Marseille	France
Isabelle Sermet-Gaudelus	Necker Hospital, Paris	France
Aurelie Tatopoulos	Nancy	France
Lutz Nährlich	Justus-Liebig-University	Germany
Barry Plant	Cork University Hospital, Cork	Ireland
Cedric Gunaratnam	Beaumont Hospital, Dublin	Ireland
Abaigeal Jackson	CF Registry of Ireland, Dublin	Ireland

Karin M. de Winter- de Groot	Wilhelmina Children's Hospital - University Medical Center Utrecht	Netherlands
Bart Luijk	University Medical Center Utrecht	Netherlands
Geertjan Wesseling	Maastricht UMC	Netherlands
Mark Allenby	Wessex Adult CF Service / University Hospital Southampton	UK
Jamie Duckers	All Wales Adult Cystic Fibrosis Service	UK
Elaine Gunn	Cystic Fibrosis Trust	UK
Andrew Jones	Royal Brompton Hospital / Imperial College London	UK
Robert Ian Ketchell	All Wales Adult Cystic Fibrosis Service	UK
Susan L. Madge	Royal Brompton Hospital	UK
Anirban Maitra	Manchester Children's Hospital	UK
Ghulam Mujtaba	Manchester Children's Hospital	UK
Helen Rodgers	Scottish Cystic Fibrosis Service	UK
Nadia Shafi	St Bartholomew's Hospital	UK
Nicholas Simmonds	Royal Brompton Hospital / Imperial College London	UK
Kevin Southern	Alder Hey Children's Hospital / University of Liverpool	UK
Danie Watson	St Bartholomew's Hospital	UK
Samar Rizvi	Cystic Fibrosis Foundation	USA
Julie Seguin	SUNY Upstate Medical University	USA
Janet Garbarz	Cook Children's Medical Center	USA
Kristen Rosamilia	Morristown Medical Center	USA
Maria Berdella	Beth Israel Medical Center	USA
Jerry A. Nick	National Jewish Health	USA
Richard Belkin	Santa Barbara Cottage Hospital	USA
Diana Gilmore	Nationwide Children's Hospital	USA
Kim McBennett	Rainbow Babies and Children's Hospital	USA

Introduction

A cluster of cases of pneumonia of unknown origin presented in Wuhan, China in early December 2019; the cause was subsequently found to be a novel coronavirus, now named SARS-CoV-2 (1). This new virus is highly transmissible and has resulted in a pandemic with over 2 million confirmed infections and more than 145,000 deaths in humans in 185 countries (2).

SARS-CoV-2 presents with variable symptoms from a mild cough, fever or flu-like illness that can resolve in a few days to a more serious pneumonia that can lead to adult respiratory distress syndrome, respiratory failure, a cytokine storm and multiple organ failure (3). The outcome appears age dependent with children frequently having minimal or mild symptoms or being asymptomatic whereas elderly people tend to have significant mortality and morbidity. Estimates of the case fatality rate (CFR) after adjustments for demography and under-ascertainment in China were 1.38% (1.23%-1.53%), with substantially higher rates in older age groups: 6.4% (5.7%-7.2%) in those over 60 years old as compared to 0.32% (0.27%-0.38%) in those under 60 years (4). Comorbidities such as diabetes mellitus, hypertension and chronic lung disease have been reported to be associated with a worse outcome and higher CFR (5). There is uncertainty regarding the true prevalence of SARS-CoV-2 and therefore the Infection Fatality Rates (IFR).

Cystic fibrosis (CF) is a multisystem genetic condition resulting in progressive chronic lung disease and multiple comorbidities including CF-related diabetes mellitus (CFRD) in up to one third of people (6), liver disease and malnutrition. In 2009-2010 the H1N1 influenza pandemic resulted in significant

morbidity in most people with CF who contracted the infection (7). Many countries have categorised people with CF as highly vulnerable to SARS-CoV-2 and have advised “shielding” or “cocooning” (staying at home at all times and avoiding face-to-face contact with anyone outside the household) to try to minimise the risk of people with CF contracting the virus. The impact of acute infection with SARS-CoV-2 in people with CF is not known (8). Here we report the clinical characteristics and outcome of 40 people across 8 countries with CF who have tested positive for SARS-CoV-2 infection.

Methods

The ‘Cystic Fibrosis Registry Global Harmonization Group’ set up extra-ordinary meetings at the outset of the SARS-CoV-2 pandemic, meeting twice monthly to share case reports and case capture methods. CF Registries in each participating country contacted all CF care centres to request data be provided. Once consensus was reached on key reporting requirements, a template was circulated to each participating country. Eight countries were able to obtain relevant ethics approvals and submit anonymised data by the cutoff date of 13 April 2020.

Anonymised data were collected, identifying neither the person with cystic fibrosis nor the treatment centre. Cases were identified as people with a definitive diagnosis of cystic fibrosis who tested positive for SARS-CoV-2 between 1 February – 13 April 2020. Cases submitted with CT scans suggestive of SARS-CoV-2 but without a positive test result for the virus were excluded. Unresolved cases were included, reported as such, and countries were asked to report zero cases as appropriate.

The information collected included sex, age, genotype group, body mass index (BMI), most recent ‘Best’ FEV1% predicted, treatment data, CF related complications, microbiology, transplant, pregnancy and current health status. Data collection is ongoing, and additional countries have committed to ongoing case capture and follow up.

Information governance

Each participating country was responsible for ensuring Research Ethics Board or institutional approvals were in place for this work. A nominated country representative collated all known cases and added them to a standardised template, accurate at 13 April 2020.

Results

A total of 8 countries are participating in the study; Australia, Canada, France, Ireland, Netherlands, New Zealand, UK and US. New Zealand reported zero cases. Table 1 reports the demographics, clinical attributes and symptom presentation of the 40 cases reported broken down into underlying CF severity according to lung function. Of the 40 cases, 31 (78%) were symptomatic for SARS-CoV-2 at presentation, with 24 (60%) having a fever. The median age was 33 years (range 15-59 years), with only one patient being under 16. 38% had CFRD and 70% were reported to have chronic bacterial pulmonary infection, of which 71% included *Pseudomonas aeruginosa* (PsA). One was pregnant and has recovered, delivering a healthy baby. Eleven reports have been from post-lung transplant patients, who are on average 6 years (range: 1-15 years) post their most recent transplant (one patient has more recently undergone a kidney transplant). Table 2 describes medical interventions and outcomes. 25 (63%) of people were started on new antibiotics: 10 acute oral antibiotics and 17 IV antibiotic therapy. Two people started on both oral and IV antibiotics.

13 (33%) patients required oxygen and 1/40 patient required invasive ventilatory support. Four (10%) were admitted to an Intensive Care Unit (ICU) for SARS-CoV-2 related reasons (two remain in ICU at time of reporting). 14 people were reported as using CFTR modulators. Of these, 1 (7%) was

admitted to ICU and 3 (21%) required oxygen. 28 of the cases have been reported as clinically recovered from SARS-CoV-2, with no deaths reported to a participating Registry.

	ALL (N=40)*		Best FEV1% predicted range						Post Tx** (N=11)	
	Media n	Rang e	Media n	Rang e	Media n	Rang e	Media n	Rang e	Media n	Rang e
Age, years	33	15-57	28	20-58	32	20-57	26	15-57	40	27-49
	N	%	N	%	N	%	N	%	N	%
Male	17	43	1	20	5	42	3	27	7	64
Homozygous F508del	17	43	3	60	3	25	4	36	6	55
Aged ≥18 years	38	95	5	100	12	100	10	91	11	100
Aged ≥40 years	13	33	2	40	3	25	2	18	6	55
	Mean	Rang e	Mean	Rang e	Mean	Rang e	Mean	Rang e	Mean	Rang e
Best FEV1% pred***	70	18-114	31	18-38	59	40-70	89	80-106	83	50-114
	Media n	Rang e	Media n	Rang e	Media n	Rang e	Media n	Rang e	Media n	Rang e
BMI, kg/m ²	22	16-34	19	18-23	22.5	16-33	23	20-34	21	16-24
	N	%	N	%	N	%	N	%	N	%
BMI >30kg/m ²	2	5	0	0	1	8	1	9	0	0
Chronic PsA	20	50	5	100	6	50	5	45	4	36
CFRD	15	38	1	20	6	50	6	55	2	18
CFTR modulator†	14	35	3	60	5	42	6	55	0	0
	N	%	N	%	N	%	N	%	N	%
Symptomatic	31	78	3	60	9	75	8	73	10	91
Fever	24	77	3	100	6	67	6	75	8	80
Dyspnea	10	32	0	0	1	11	5	63	4	40
Altered cough	15	48	2	67	5	56	3	38	5	50

Table 1 – Demographics, clinical attributes and symptom presentation

	No transplant		Post-transplant**		All	
	N = 29	%	N = 11	%	N=40	%
Interventions						
Intensive care (SARS-COV-2 related)	1	3	3	27	4	10
New supplemental oxygen	7	24	6	55	13	33
Invasive ventilation	0	0	1	9	1	3
Outcomes						
Clinically recovered	22	76	6	55	28	70
Unresolved at time of report	7	24	5	45	12	30
Died	0	0	0	0	0	0

Table 2 - Interventions and outcomes

* One patient, who has not received a lung transplant, is not able to perform spirometry testing, so is excluded from the 'Best FEV1% predicted range' counts. This patient only appears in the 'ALL' counts.

** Lung, Lung and Liver or Lung and Kidney (consecutive) transplants (Tx)

*** Best FEV1% predicted. Countries provided the most recent available pre-infection best FEV1% predicted. Measurements are from 2019 or 2020

† Ivacaftor, lumacaftor/ivacaftor, tezacaftor/ivacaftor, or elaxacaftor/tezacaftor/ivacaftor (9)

Discussion

These early data suggest that the course of disease in CF may not be as severe as expected from initial data from patients with other underlying lung diseases. This CF population infected with SARS-CoV-2 are heterogeneous, with a wide range of lung function, complications, chronic respiratory infections. No apparent themes or specific risk factors for the contraction or severity of SARS-CoV-2 are detectable at this stage. No fatalities have been reported in this series, despite the presence of low lung function, lung transplantation listing or receipt, pregnancy and serious comorbidities amongst the cohort. The only patient to date needing invasive ventilation associated with SARS-CoV-2 has been in an individual who was post-transplant.

The median age of this cohort at 33 years is higher than the general CF population, for comparison the median age in the UK CF population was 20 in 2018 (6). This is due to the low number of reported cases in the pediatric population. This adult predominance in CF reflects the picture for SARS-CoV-2 in the general population (8). 43% of this cohort are male, unlike current reports of 60-75% of admitted patients in the general population being male (10). The median BMI of adults aged 20 or over in the UK CF population is 22.5kg/m² (6), it was 22kg/m² in this SARS-CoV-2 cohort.

Eleven (28%) people with CF in the cohort have had a lung transplant. Post-transplant patients with CF may be over-represented in this cohort as a result of transplant centre protocols to investigate new respiratory symptoms, a change in lung function and a new temperature in post-transplant patients. This is borne out by the high proportion (91%) of post-transplant patients tested due to presentation of symptoms rather than routine screening.

A high proportion of this SARS-CoV-2 cohort received acute oral or IV antibiotics during their infection. Antibiotics are not known to be effective against SARS-CoV-2 and were administered to treat the underlying chronic respiratory infection or exacerbation.

To date, incidence of SARS-CoV-2 amongst the CF population (0.07%) appears to be lower than the average derived on 13 April from the general populations of participating countries (0.15%) (11), all countries within this study are at different stages of the pandemic, as such general population rates will differ. The apparent lower rate in CF may be partly attributable to the lower average age of the CF population, lags in reporting. It may also be due to people with CF adopting “shielding”, “protective self-isolation” or “cocooning” (staying at home at all times and avoiding face-to-face contact with anyone outside the household) earlier and being more effective at mitigating the risk of cross infection than the general public. It has also been suggested that the chronic respiratory disease population is ‘primed’ for reducing risk of exposure due to learned behaviors or standard of care therapies (12).

Limitations

The SARS-CoV-2 pandemic is a rapidly evolving global situation with participating countries at different phases of the epidemic. Not all countries in Europe were able to participate in this study, notably some countries with substantial experience of SARS-CoV-2. Reasons for non-participation were lack of a national data collection infrastructure, reporting lag, or delays gaining ethics approvals for data sharing. For participating countries, this method of case capture is not exhaustive and may not identify all cases. This may be due to inability to obtain all required data or test results

prior to the deadline, or delayed reporting by clinical teams to their respective Registry. Cases where CT scans were highly suggestive of SARS-CoV-2 but the patient had not tested positive for the virus were excluded from this study. In some cases, multiple tests had been performed, but reports of 20-30% false-negative SARS-CoV-2 test rates should be noted (13). There are varying testing protocols across the participating countries and ascertaining the number of tests conducted in the CF population is hard, meaning the denominator of people with CF who have been exposed to SARS-CoV-2 is not known.

Caution must be exercised when interpreting these early results, and the limitations of this study born in mind. We cannot combine the national case series to estimated mortality because of heterogenous testing policies and reporting processes across different countries.

Conclusions

The outcomes of this early cases of SARS-CoV-2 in 40 people with CF, reported across 8 countries have been better than initially predicted based on previous respiratory pandemics. However, the medium and long-term impact of SARS-CoV-2 amongst infected patients is still not known.

The absence of mortality amongst the population may be linked to the relatively low incidence of SARS-CoV-2 amongst the CF population. This could be attributed to effective shielding techniques, but it is too early to state this with confidence.

Conflict of interest statement

The authors have no conflicts of interest to declare relating to this work.

References

1. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 2020 Feb 28;NEJMoa2002032. doi: 10.1056/NEJMoa2002032. Epub ahead of print. PMID: 32109013; PMCID: PMC7092819.
2. <https://coronavirus.jhu.edu/map.html> (Accessed: 16 April 2020)
3. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020 Feb 15;395(10223):507-513. doi: 10.1016/S0140-6736(20)30211-7. Epub 2020 Jan 30. PMID: 32007143; PMCID: PMC7135076.
4. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis*. 2020 Mar 30:S1473-3099(20)30243-7. doi: 10.1016/S1473-3099(20)30243-7. Epub ahead of print. PMID: 32240634.
5. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med*. 2020 Mar 13:e200994. doi: 10.1001/jamainternmed.2020.0994. Epub ahead of print. PMID: 32167524; PMCID: PMC7070509.
6. UK Cystic Fibrosis Registry Annual Data Report 2018, Cystic Fibrosis Trust
7. Viviani L, Assael BM, Kerem E; ECFS (A) H1N1 study group. Impact of the A (H1N1) pandemic influenza (season 2009-2010) on patients with cystic fibrosis. *J Cyst Fibros*. 2011 Sep;10(5):370-6. doi: 10.1016/j.jcf.2011.06.004. Epub 2011 Jul 12. PMID: 21752728.
8. Colombo C, Burgel P-R, Gartner S, van Koningsbruggen-Rietschel S, Naerlich L, Sermet-Gaudelus I et al. Impact of COVID-19 on people with cystic fibrosis. *Lancet Respir Med* 2020 Apr 15: S2213-2600(20)30177-6. doi: 10.1016/S2213-2600(20)30177-6. Epub ahead of print. PMID: 32304639.
9. Heijerman HGM, McKone EF, Downey D G, Braeckel EV, Rowe SM, Tullis E et al. Efficacy and safety of the elxacaftor plus tezacaftor plus ivacaftor combination regimen in people with cystic fibrosis homozygous for the *F508del* mutation: a double-blind, randomised, phase 3 trial. *Lancet* 2019 Nov 23;394(10212):1940-1948. doi: 10.1016/S0140-6736(19)32597-8. PMID: 31679946.
10. Yang X, Xu J, Shu H, Xia J, Liu H, Wu Y et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centred, retrospective, observational study. *Lancet Respir Med* 2020 Feb 21: S2213-2600(20)30079-5. doi: 10.1016/S2213-2600(20)30079-5. Epub ahead of print. PMID: 32105632.
11. Johns Hopkins University (Accessed: 13 April 2020) and World Bank 2018 population estimates (Accessed: 13 April 2020).
12. Halpin DM, Farber R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection?. *Lancet Respir Med* 2020 Apr 3: S2213-2600(20)30167-3. doi: 10.1016/S2213-2600(20)30167-3. Epub ahead of print. PMID: 32251625.
13. Li D, Wang D, Dong J, Wang N, Huang H, Xu H et al. False-negative results of real-time Reverse-Transcriptase Polymerase Chain Reaction for severe acute respiratory syndrome Coronavirus 2:

Role of deep-learning-based CT diagnosis and insights from two cases. Korean J Radiol 2020 Apr;21(4):505-508. English. doi: 10.3348/kjr.2020.0146. PMID: 32174053.

Credit author statement

Rebecca Cosgriff: Conceptualization, Methodology, Formal analysis, Investigation, Data Curation, Writing – Original Draft, Writing – Review & Editing, Project administration **Susannah Ahern:** Investigation, Data Curation, Writing – Review & Editing **Scott C. Bell:** Writing – Review & Editing **Keith Brownlee:** Conceptualization, Writing – Original Draft, Writing – Review & Editing **Pierre-Régis Burgel:** Investigation, Data Curation, Writing – Review & Editing **Cass Byrnes:** Conceptualization, Investigation **Harriet Corvol:** Investigation, Data Curation, Writing – Review & Editing **Stephanie Y. Cheng:** Conceptualization, Investigation, Data Curation, Writing – Review & Editing **Alexander Elbert:** Conceptualization, Investigation, Data Curation, Writing – Review & Editing **Albert Faro:** Conceptualization, Writing – Review & Editing **Christopher H. Goss:** Conceptualization, Writing – Review & Editing **Vincent Gulmans:** Conceptualization, Investigation, Data Curation, Writing – Review & Editing **Bruce C. Marshall:** Conceptualization, Writing – Review & Editing **Edward McKone:** Conceptualization, Investigation, Data Curation, Writing – Review & Editing **Peter G. Middleton:** Investigation, Data Curation, Writing – Review & Editing **Rasa Ruseckaite:** Investigation, Data Curation, Writing – Review & Editing **Anne L. Stephenson:** Conceptualization, Writing – Review & Editing **Siobhán B Carr:** Conceptualization, Validation, Writing – Original Draft, Writing – Review & Editing, Supervision